

# Combined Effect of Cigarette Smoke and Mineral Fibers on the Gene Expression of Cytokine mRNA

Yasuo Morimoto,<sup>1</sup> Tohru Tsuda,<sup>1</sup> Hajime Hori,<sup>2</sup> Hiroshi Yamato,<sup>3</sup> Akira Ohgami,<sup>3</sup> Toshiaki Higashi,<sup>4</sup> Nobuhiko Nagata,<sup>5</sup> Masamitsu Kido,<sup>5</sup> and Isamu Tanaka<sup>3</sup>

<sup>1</sup>Department of Occupational Pneumology, Institute of Industrial Ecological Sciences; <sup>2</sup>School of Health Sciences; <sup>3</sup>Department of Environmental Health Engineering, Institute of Industrial Ecological Sciences; <sup>4</sup>Department of Work Systems and Health, Institute of Industrial Ecological Sciences; and <sup>5</sup>Department of Respiratory Diseases, University of Occupational and Environmental Health, Japan, Kitayushu, Japan

To investigate which parameters are stimulated by mineral fibers and whether cigarette smoke enhanced a fiber-induced response, we examined the level of cytokine mRNA from alveolar macrophages (AMs) and lungs of rats exposed to mineral fibers and cigarette smoke *in vivo*. Male Wistar rats were given a single intratracheal instillation of 2 mg of Union Internationale Contre le Cancer chrysotile or refractory ceramic fiber (RF1). The animals then inhaled a side stream of smoke 5 days per week for 4 weeks. The expression of manganese superoxide dismutase, inducible nitric oxide synthase (iNOS), basic fibroblast growth factor (bFGF), interleukin-1 $\alpha$  (IL-1 $\alpha$ ), interleukin-6 (IL-6), and tumor necrosis factor- $\alpha$  (TNF $\alpha$ ) mRNA from lipopolysaccharide-stimulated AMs and lungs of rats exposed to mineral fibers and/or cigarette smoke were assessed using semiquantitative reverse-transcriptase polymerase chain reaction. Exposure only to cigarette smoke increased in IL-1 $\alpha$  mRNA levels in AMs. Chrysotile stimulated the expression of IL-1 $\alpha$ , TNF $\alpha$ , and IL-6 in AMs, and the expression of bFGF in lungs. RF1 resulted in increased expression of IL-1 $\alpha$  and TNF $\alpha$  in AMs. Cigarette smoke stimulated the gene expression of iNOS in AMs and IL-6 and bFGF in lungs treated with chrysotile; IL-1 $\alpha$  in AMs and bFGF in lungs did the same in lungs with RF1. Among these cytokines, message levels of IL-1 $\alpha$ , iNOS, and bFGF were increased in rats stimulated with mineral fibers, and the stimulating effects of mineral fibers were enhanced by cigarette smoke. Therefore, IL-1 $\alpha$ , iNOS, and bFGF would be the possible parameters of the lung remodeling induced by mineral fibers. **Key words:** alveolar macrophage, bFGF, ceramic fiber, chrysotile, IL-1 $\alpha$ , IL-6, iNOS, lung, Mn-SOD, TNF. *Environ Health Perspect* 107:495–500 (1999). [Online 7 May 1999]

<http://ehpnet1.niehs.nih.gov/docs/1999/107p495-500morimoto/abstract.html>

Occupational and environmental exposure to inhaled asbestos dust causes pulmonary fibrosis, bronchogenic carcinoma, and pleural mesothelioma (1). Recently, various types of man-made mineral fibers have been developed as substitutes for asbestos; the demand for these products is increasing. Some of these fibers are thought to possess the same adverse biological effects as asbestos because of their similar physiochemical properties (2). Furthermore, epidemiological and clinical studies have indicated that asbestos workers who smoked cigarettes had a higher incidence of asbestos-induced disease than those who did not (3–4).

In asbestos-induced disease, many genes are thought to contribute to lung remodeling, such as fibrosis and carcinoma (5–6). Asbestos fibers deposited in the lung lead to an activation of alveolar macrophages (AMs) (5). AMs and parenchymal cells release tumor necrosis factor- $\alpha$  (TNF $\alpha$ ), interleukin-1 $\alpha$  (IL-1 $\alpha$ ), interleukin-6 (IL-6), basic fibroblast growth factor (bFGF), and other cytokines that augment the process of inflammation (5–6). The release of oxidants from these cells may lead to lung injury and manganese superoxide dismutase (Mn-SOD) and inducible nitric oxide synthase (iNOS) corresponding

to released free radical formation (6). Some growth factors signal interstitial fibroblasts to replicate and modulate their production of connective tissue proteins (5). The accumulation of inflammatory cells and fibroblasts and the development of connective tissue matrices causes lung injury and leads to fibrosis and carcinoma (5–6).

A reverse transcriptase-polymerase chain reaction (RT-PCR) used in this study has a number of advantages (7): *a*) its exquisite sensitivity gives it the ability to detect rare mRNA in small numbers of cells in a semiquantitative manner; *b*) unlike the protein assay system, the same methodology can be applied to analyze the expression of many genes; and *c*) cDNA can be used for future studies.

To estimate which parameters are more sensitive and specific for lung remodeling induced by mineral fibers, we used RT-PCR to examine cytokine mRNA levels from AMs and lungs of rats exposed to mineral fibers and/or cigarette smoke *in vivo*.

## Materials and Methods

**Fibers.** Fibers used in this study were Union Internationale Contre le Cancer chrysotile and alumina silicate ceramic fibers (RF1).

For chrysotile, the geometric mean diameter [standard deviation (SD)] and the geometric mean length were 0.085 (1.4)  $\mu$ m and 0.7 (0.19)  $\mu$ m, respectively (2). For RF1, the geometric mean diameter (SD) and the geometric mean length were 0.77 (2.53)  $\mu$ m and 12.0 (2.36)  $\mu$ m, respectively. RF1 was provided by the Japan Fibrous Material Research Association (Tokyo, Japan).

**Experimental design.** Male Wistar rats (10 weeks old) were used in this study, and 30 rats were divided into six groups (saline, chrysotile, RF1, cigarette smoke, chrysotile + cigarette smoke, and RF1 + cigarette smoke). Either saline, chrysotile suspension, or RF1 suspension (2 mg/0.2 mL) was administered to the rats intratracheally. The rats were housed in an exposure chamber and were exposed to the smoke of 20 cigarettes (Japan Tobacco, Tokyo, Japan) for 4 hr/day, 5 days/week, for 4 weeks. The mass concentration of cigarette smoke was 10 mg/m<sup>3</sup> and was measured gravimetrically at daily intervals by the suction of air through a glass filter. The CO concentration was 79 ppm, which is below the concentration that induces specific biological effects. The chamber volume was 100 L and the flow rate in the chamber was 50 L/min.

**Bronchoalveolar lavage and preparation.** After exposure, each animal was anesthetized by an intraperitoneal injection of phenobarbital. Bronchoalveolar lavage (BAL) was performed on the left lung. Cells recovered by BAL were suspended in RPMI-1640 medium, then placed on tissue culture plates and allowed to attach for 1 hr at 37°C (8). Nonadherent cells were removed by one cycle of washing with RPMI, and adherent cells were supplied with an RPMI-1640 medium

Address correspondence to Y. Morimoto, Department of Occupational Pneumology, Institute of Industrial and Ecological Sciences, University of Occupational and Environmental Health, Japan, 1-1 Iseigaoka, Yahatanishi, Kitakyushu, Japan. Telephone: +81 93 691 7466. Fax: +81 93 691 4284. E-mail: yasuo@med.uoeh-u.ac.jp

We thank M. Tsunoda and M. Hirohashi for assistance with some of the experiments. This work was supported in part by the Ministry of Education and Science in Japan, the Fukuoka Cancer Society, and the Uehara Memorial Foundation.

Received 17 December 1997; accepted 8 February 1999.

containing 10% fetal bovine serum. They were adjusted to a final concentration of  $1 \times 10^5$ /mL and stimulated with 10  $\mu$ g/mL lipopolysaccharide (Sigma, St. Louis, MO). The AMs were cultured for 2 and 6 hr on cell culture plates at 37°C in the CO<sub>2</sub> incubator. After incubation, mRNA in AMs was extracted using a Quick Prep kit (Pharmacia Biotech, Uppsala, Sweden). Total RNA from the lung was prepared in the presence of guanidium thiocyanate (9).

**Preparation of RNA, cDNA synthesis, and PCR.** Total RNA (0.5  $\mu$ g) was used for the synthesis of single-strand cDNA using Moloney murine leukemia virus-derived reverse transcriptase (Perkin Elmer, Norwalk, CT). An equal amount of cDNA from each sample, standardized to give identical signals on gel following amplification with  $\beta$ -actin primer, was amplified by specific primers for each gene (Table 1) (10–16). The amplification was performed using a Thermocycler (Astech, Japan) under the following conditions: denaturation at 94°C for 45 sec, annealing at 60°C for 45 sec, and extension at 72°C for 2 min for cytokines and  $\beta$ -actin genes.

Detection of the fragments amplified by PCR was made by electrophoresis on a 2% agarose gel. PCR products were resolved using gel electrophoresis and visualized by ethidium bromide staining. The gel was photographed with Polaroid Type 665 positive/negative film (Polaroid Corp., Cambridge, MA) under ultraviolet light at identical exposure and development times. The bands from the positive film were scanned, and the density of each PCR product was measured using National Institute of Health image 1.55 software (provided by Wane Rasband, National Institutes of Health, Bethesda, MD). The ratio of each specific gene product to the  $\beta$ -actin product was used for the analysis.

**Quantitative analysis.** To quantify the transcriptional level of cytokine mRNA, the following analysis was used: IL-1 $\alpha$  samples were coamplified with  $\beta$ -actin-specific primers as an internal standard (Figure 1). Aliquots containing cDNA were subjected to 26, 29, 32, 35, or 38 cycles of amplification under identical conditions as in protocols. The amounts of density recovered from the excised bands were plotted as a function of the number of cycles. The rates of amplification were exponential between 29 and 35 cycles for both templates. The number of thermocycles used allowed quantitation without saturation. The data were normalized to represent equivalent RNA loading based on the density of  $\beta$ -actin at the appropriate cycle of both genes (7).

We compared the data of 2- and 6-hr incubation of cytokines in AMs and presented the one that showed more gene expression.

**Statistical analysis.** Values were expressed as the mean  $\pm$  1 SD. The difference between values was assessed using a Mann-Whitney U-test.

**Histopathology.** Lungs were fixed in 10% buffered formalin. Embedded samples were then sectioned and stained with hematoxylin and eosin.

## Results

**Bronchoalveolar lavage.** An increased number of total cells was recovered from rats exposed to mineral fibers and/or cigarette smoke than from saline-exposed rats (cigarette  $6.4 \pm 3.4 \times 10^5$  cells; chrysotile  $5.6 \pm 1.9 \times 10^5$  cells; RF1  $5.5 \pm 1.9 \times 10^5$  cells; chrysotile + cigarette smoke  $5.7 \pm 1.5 \times 10^5$  cells; RF1 + cigarette smoke  $9.1 \pm 1.6 \times 10^5$  cells; control  $4.4 \pm 1.2 \times 10^5$  cells; Table 2). AMs accounted for >95% of the cells in all groups.


**Expression of IL-1 $\alpha$ , TNF $\alpha$ , and IL-6 mRNA.** In comparison with the control (saline) group, levels of IL-1 $\alpha$  mRNA in AMs increased significantly in groups exposed to cigarette smoke, chrysotile, chrysotile + cigarette smoke, RF1, and RF1 + cigarette smoke (Figure 2A). IL-1 $\alpha$  mRNA in the group exposed to RF1 + cigarette smoke was maximally expressed in all groups, but it was not significant compared to cigarette smoke- or RF1-exposed group. Elevation in the gene expression of IL-1 $\alpha$  mRNA was not observed in rat lungs exposed to mineral fibers and/or cigarette smoke (Figure 2B).

A significant increase in TNF $\alpha$  mRNA in AMs was obtained in chrysotile- or

**Table 1.** Oligonucleotides of primers of seven target genes.

mRNA species	mRNA	Cycle of PCR product, bp		
		Lung	AMs	bp
TNF $\alpha$	(sense) 5'-TACTGAACCTTCGGGGTGATTGGTCC	30	32	295
	(antisense) 5'-CAGCCTTGTCCTTGAAGAGAACC			
IL-1 $\alpha$	(sense) 5'-CTAAGAACTACTTCACATCCGCAGC	38	32	623
	(antisense) 5'-CTGGAATAAAACCCACTGAGGTAGG			
IL-6	(sense) 5'-CAAGAGACTTCAGCCAGTTGC	40	40	614
	(antisense) 5'-TTGCCGAGTAGACCTCATAGTGACC			
iNOS	(sense) 5'-CAAGAACGTGTTCAACCATGA	37	40	696
	(antisense) 5'-AAAGCAGGGCACTGGGTCTT			
Mn-SOD	(sense) 5'-GCGACCTACGTGAACAATCTGAACG	28	40	383
	(antisense) 5'-TCAATCCCCAGCAGTGGAATAAGGC			
bFGF	(sense) 5'-CAAGCAGAAGAGAGAGGAGTT	36	40	260
	(antisense) 5'-TCAGCTCTTAGCAGACATTG			
$\beta$ -Actin	(sense) 5'-ATCATGTTTGAGACCTTCAACACC	26–28	32–34	357
	(antisense) 5'-TAGCTCTTCTCCAGGGAGG			

Abbreviations: AMs, alveolar macrophages; bFGF, basic fibroblast growth factor; IL-1 $\alpha$ , interleukin-1 $\alpha$ ; IL-6, interleukin-6; iNOS, inducible nitric oxide synthase; Mn-SOD, manganese superoxide dismutase; PCR, polymerase chain reaction; TNF $\alpha$ , tumor necrosis factor  $\alpha$ .



**Figure 1.** Gene expression of interleukin-1 $\alpha$  (IL-1 $\alpha$ ) mRNA in alveolar macrophages of rats exposed to chrysotile or saline in a cycle-dependent fashion. (A) Exposure to saline. (B) Exposure to chrysotile.




RF1-exposed rats as compared to control rats, but not in cigarette smoke-exposed rats (Figure 3A). Mineral fibers and cigarette smoke did not have combined effects on message levels of TNF $\alpha$  in AMs. TNF $\alpha$  mRNA in the lung was increased in chrysotile-exposed rats. Similar to chrysotile, RF1 tended to increase in the gene expression of TNF $\alpha$  mRNA; however, there were no significant differences in the levels of TNF $\alpha$  mRNA when compared to control rats (Figure 3B).

Chrysotile-only exposure induced increases in IL-6 mRNA levels in AMs (Figure 4A). Elevation of IL-6 mRNA was not increased in rat AMs exposed to RF1 and/or cigarette smoke. Exposure to cigarette smoke or mineral fibers did not induce an

**Table 2.** Total cell counts of bronchoalveolar lavage fluid in rats exposed to mineral fibers and/or cigarette smoke *in vivo*.

	Total cell counts ( $\times 10^5$ )
Control	4.4 (1.2)
Cigarette	6.4 (3.4)
Chrysotile	5.6 (1.9)
RF1	5.5 (1.9)
Chrysotile + cigarette smoke	5.7 (1.5)
RF1 + cigarette smoke	9.1 (1.6)

RF1, refractory ceramic fiber.



**Figure 2.** Effects of mineral fibers and cigarette smoke on the expression of interleukin-1 $\alpha$  (IL-1 $\alpha$ ) mRNA. (A) Alveolar macrophages incubated for 2 hr. (B) Lung. Abbreviations: NS, not significant; RF1, refractory ceramic fiber. Values are significantly greater than control (saline). Results are expressed in ratio of IL-1 $\alpha$  to  $\beta$ -actin (mean  $\pm$  standard error).


\* $p < 0.05$  significantly different from a control (saline-exposed) group.

increase in IL-6 levels in the lung; however, exposure to chrysotile and cigarette smoke had a synergistic effect on the expression of IL-6 mRNA (Figure 4B).

**Expression of iNOS and Mn-SOD mRNA.** In rats exposed to cigarette smoke or mineral fibers, mRNA levels for iNOS in AMs were not increased (Figure 5A). On the other hand, mRNA levels for iNOS


among AMs from rats treated with mineral fibers and cigarette smoke increased significantly when compared with control rats.

Levels of iNOS mRNA declined markedly in cigarette smoke-exposed rat lungs (Figure 5B). Exposure to mineral fibers tended to induce an increase in iNOS mRNA; however, differences between mineral fibers and control were not significant.




**Figure 3.** Effects of mineral fibers and cigarette smoke on the expression of tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) mRNA. (A) Alveolar macrophages incubated for 2 hr. (B) Lung. Abbreviations: NS, not significant; RF1, refractory ceramic fiber. Results are expressed in ratio of TNF $\alpha$  to  $\beta$ -actin (mean  $\pm$  standard error).

\* $p < 0.05$  and \*\* $p < 0.01$  significantly different from a control (saline-exposed) group.



**Figure 4.** Effects of mineral fibers and cigarette smoke on the expression of interleukin-6 (IL-6) mRNA. (A) Alveolar macrophages incubated for 6 hr. (B) Lung. Abbreviations: NS, not significant; RF1, refractory ceramic fiber. Results are expressed in ratio of IL-6 to  $\beta$ -actin (mean  $\pm$  standard error).


\* $p < 0.05$  significantly different from a control (saline-exposed) group.




**Figure 5.** Effects of mineral fibers and cigarette smoke on the expression of inducible nitric oxide synthase (iNOS) mRNA. (A) Alveolar macrophages incubated for 2 hr. (B) Lung. Abbreviations: NS, not significant; RF1, refractory ceramic fiber. Results are expressed in ratio of iNOS to  $\beta$ -actin (mean  $\pm$  standard error).

\* $p < 0.05$  and \*\* $p < 0.01$  significantly different from a control (saline-exposed) group.






**Figure 6.** Effects of mineral fibers and cigarette smoke on the expression of manganese superoxide dismutase (Mn-SOD) mRNA. (A) Alveolar macrophages incubated for 6 hr. (B) Lung. Abbreviations: NS, not significant; RF1, refractory ceramic fiber. Results are expressed in ratio of Mn-SOD to  $\beta$ -actin (mean  $\pm$  standard error).



**Figure 8.** Lung sections in rats exposed to mineral fibers and/or cigarette smoke. (A) Saline, magnification  $\times 100$ . (B) Cigarette smoke, magnification  $\times 100$ . (C) Chrysotile, magnification  $\times 400$ . (D) refractory ceramic fiber (RF1), magnification  $\times 100$ . (E) Chrysotile + cigarette smoke, magnification  $\times 400$ . (F) RF1 + cigarette smoke, magnification  $\times 100$ .



**Figure 7.** Effects of mineral fibers and cigarette smoke on the expression of basic fibroblast growth factor (bFGF) mRNA. (A) Alveolar macrophages incubated for 2 and 6 hr. (B) Lung. Abbreviations: NS, not significant; RF1, refractory ceramic fiber. Results are expressed in ratio of bFGF to  $\beta$ -actin (mean  $\pm$  standard error).

\* $p < 0.05$  and \*\* $p < 0.01$  significantly different from a control (saline-exposed) group.


No significant changes in levels of Mn-SOD mRNA in AMs and lungs exposed to mineral fibers and/or cigarette smoke was observed (Figure 6).

**Expression of bFGF mRNA.** The message of bFGF in AMs was not observed in all groups, although the cycle of PCR was up to 40 (Figure 7A). Chrysotile induced an increase in bFGF mRNA in the lungs, but cigarette smoke or RF1 did not. The combined effect of mineral fibers and cigarette smoke on the message level of bFGF was significant (Figure 7B).

**Pathological findings.** Foreign body giant cells and mononuclear cells around bronchioles were observed in groups exposed to chrysotile and chrysotile + cigarette smoke (Figure 8). In the other four groups, definite pathological changes were not observed.

## Discussion

We set the recovery time after the intratracheal instillation of mineral fibers at 4 weeks. Unlike the inhalation findings, some reports showed that nonspecific responses at the acute phase were seen after the intratracheal injection. Some investigators (17,18)



**Figure 9.** Ethidium bromide staining of PCR products separated in 2% agarose gel. Abbreviations: bFGF, basic fibroblast growth factor; IL-1 $\alpha$ , interleukin-1 $\alpha$ ; IL-6, interleukin-6; M, DNA marker ( $\phi$ X174/*Hae*III); Mn-SOD, manganese superoxide dismutase; iNOS, inducible nitric oxide synthase; RCF, refractory ceramic fiber; TNF $\alpha$ , tumor necrosis factor  $\alpha$ .

showed that saline injection in BAL caused a neutrophil recruitment by the lower respiratory tract in 72 hr, and that an increase in neutrophils and AMs recovered from bronchoalveolar lavage fluid were seen in the first week after injecting water alone. We previously reported that inhalation for one day of cigarette smoke transiently enhanced the ability of mineral fibers to stimulate AMs to produce TNF in rats (19), and later reported

that no specific changes were found for AMs in rats with cigarette smoke inhalation for 4 weeks (20). To avoid the initial influence of acute response, we exposed rats to cigarette smoke for 4 weeks in this experiment. The time line for gene expression of cytokines in AMs and lungs following initial asbestos exposure is now under investigation.

Many previous reports examined protein level or gene expression of proinflammatory

cytokines and growth factors that were stimulated by mineral fibers in lungs or in AMs (5,6,19–21). However, the mineral fibers used for experimental studies are not always the same among laboratories. The physicochemical properties of mineral fibers affect biological response in cells and lungs. For instance, the size of mineral fibers causes different responses even if they are the same fibers. We previously reported a difference in the production of TNF $\alpha$  in AMs treated with long and short mineral fibers (22). Numerous *in vitro* studies have also shown that long fibers induced greater cytotoxicities than short fibers *in vitro* (2,23,24). In this experiment each cytokine was examined from the same samples using RT-PCR under the identical experimental process, and it is useful to estimate the sensitivity and selectivity of parameters in lung remodeling induced by mineral fibers. In this experiment, these parameters were examined on the following points: whether the gene expression of cytokine mRNA is induced by mineral fibers and whether mRNA levels for cytokines are additively or synergistically increased by combined exposure to cigarette smoke and mineral fibers.

Mineral fibers increased in gene expression of TNF $\alpha$ , IL-1, IL-6, and iNOS mRNA in AMs and bFGF mRNA in lungs. These proinflammatory cytokine mRNA or proteins have been detected in the AMs or lungs of patients with idiopathic pulmonary fibrosis (25) and in animal lungs with pulmonary fibrosis (26). Blackford et al. (27) reported that intratracheal instillation of silica up regulated the gene expression of iNOS mRNA in AMs. These data were accordance with our data. However, mRNA expression of Mn-SOD was not increased in AMs and lung tissue from rats exposed to mineral fibers. Quinlan et al. (28) reported that the message level of Mn-SOD in rat lungs inhaling crocidolite peaked at 3–9 days and decreased toward control level thereafter. In our experiments, levels of Mn-SOD mRNA might return from peak to normal because recovery time was 4 weeks.

Cigarette smoke is thought to increase the incidence of asbestos-induced disease (3,4). It is generally accepted that smoking is associated with a higher risk of carcinoma of the lung (3); however, its association with asbestosis is controversial, and a majority of studies showed a positive interaction between smoking and asbestos (4). Studies suggest that smoking and asbestos act in combination in the development of fibrosis. In our experiments, cigarette smoke and mineral fibers have combined effects of gene expression of IL-1 $\alpha$ , iNOS, bFGF, and IL-6 mRNA (Figure 9). There



have been few experimental studies to investigate the combined effects of cigarette smoke and mineral fibers on the gene expression of proinflammatory cytokines. We previously reported that the cigarette smoke and mineral fibers had synergistic effects on matrix metalloproteinases (MMPs) in rat lungs (29). IL-1 up regulates the production of MMPs (30), and the increase in gene expression of MMPs in rats lungs exposed by two agents may be affected by IL-1. Jackson et al. (31) reported that cigarette smoke and asbestos increased DNA damage synergistically and suggested that this synergism might involve hydroxyl radical production. Free radical formation including nitric oxide radical and peroxynitrite anion through iNOS (27) might be related to lung injury induced by asbestos and cigarette smoke. The gene expression of proinflammatory cytokines, iNOS, and bFGF might be representative cytokines of inflammation, pulmonary damage by oxidants, and a neovascularization, respectively (4,27,32). Therefore, a change in gene expression of these factors, which play a key role in the inflammatory process, suggested that these factors might contribute to lung remodeling induced by cigarette smoke and mineral fibers. The lungs in rats exposed to chrysotile or chrysotile + cigarette smoke showed inflammatory cells around the bronchioles without a fibrotic form. These pathological findings may support that two agents have a combined effect on cytokines which are related to inflammation or lung injury.

Taken together, IL-1 $\alpha$ , iNOS, and bFGF mRNA were up regulated by exposure to mineral fibers, and were affected by exposure to mineral fibers and cigarette smoke in combination. Therefore, these factors were thought to be possible parameters for a risk assessment of mineral fibers.

## Summary

We examined cytokine mRNA levels from AMs and the lungs of rats exposed to mineral fibers and/or cigarette smoke *in vivo*. Message levels of IL-1 $\alpha$ , iNOS, and bFGF were increased by exposure to mineral fibers and enhanced by combined exposure to mineral fibers and cigarette smoke. These data suggested that IL-1 $\alpha$ , iNOS, and bFGF would be the possible parameters of the lung remodeling induced by mineral fibers.

## REFERENCES AND NOTES

- Becklake MR. Asbestos-related diseases of the lung and other organs: their epidemiology and implications for clinical practice. *Am Rev Respir Dis* 114:187-227 (1976).
- Fujino A, Hori H, Higashi T, Morimoto Y, Tanaka I, Kaji H. *In-vitro* biological study to evaluate the toxic potentials of fibrous materials. *Int J Occup Environ Health* 1:21-28 (1995).
- Hammond EC, Selikoff IJ, Seidman H. Asbestos exposure, cigarette smoking and death rates. *Ann NY Acad Sci* 330:473-490 (1979).
- Weiss W. Cigarette smoke, asbestos, and small irregular opacities. *Am Rev Respir Dis* 130:293-301 (1984).
- Warheit DB, Gavett SH. Current concepts in the pathogenesis of particulate-induced lung injury. In: *Fiber Toxicology* (Warheit DB, ed). San Diego, CA:Academic Press, 1993:305-322.
- Mossman BT, Churg A. Mechanisms in the pathogenesis of asbestosis and silicosis. *Am J Respir Crit Care Med* 157:1666-1680 (1998).
- Wang AM, Mark DF. Quantitative PCR. In: *PCR Protocols: A Guide to Methods and Applications* (Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds). San Diego, CA:Academic Press, 1990:70-75.
- Jordana M, Richards C, Irving LB, Gaudie J. Spontaneous *in vitro* release of alveolar macrophage cytokines after the intratracheal instillation of bleomycin in rats. *Am Rev Respir Dis* 137:1135-1140 (1988).
- Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162:156-159 (1987).
- Shirai T, Shimizu N, Horiguchi S, Ito H. Cloning and expression in *Escherichia coli* of the gene for rat tumor necrosis factor. *Agric Biol Chem* 53:1733-1736 (1989).
- Nishida T, Nishino N, Tanano M, Sekiguchi Y, Kawai K, Mizuno K, Nakai S, Masui Y, Hirai Y. Molecular cloning and expression of rat interleukin-1 $\alpha$  cDNA. *J Biochem* 105:351-357 (1989).
- Northemann W, Braciak TA, Hattori M, Lee F, Fey GH. Structure of the rat interleukin-6 gene and its expression in macrophage-derived cells. *J Biol Chem* 264:16072-16082 (1989).
- Kosuga K, Yui Y, Hattori R, Sase K, Eizawa H, Aoyama T, Inoue R, Sasayama S. Cloning of an inducible nitric oxide synthase from rat polymorphonuclear neutrophils. *Endothelium* 2:217-221 (1994).
- Shimasaki S, Emoto N, Koba A, Mercado M, Shibata F, Cooksey K, Baird A, Ling N. Complementary DNA cloning and sequencing of rat ovarian basic fibroblast growth factor and tissue distribution study of its mRNA. *Biochem Biophys Res Commun* 157:256-263 (1988).
- Ho YS, Crapo JD. Nucleotide sequences of cDNAs coding for rat manganese-containing superoxide dismutase. *Nucleic Acids Res* 15(23):10070 (1987).
- Nudel U, Zakut R, Shani M, Neuman S, Levy Z, Yaffe D. The nucleotide sequence of the rat cytoplasmic beta-actin gene. *Nucleic Acids Res* 11:1759-1771 (1983).
- Von Essen SG, Robbins RA, Spurzem JR, Thompson AB, McGranaghan SS. Bronchoscopy with bronchoalveolar lavage causes neutrophil recruitment to the lower respiratory tract. *Am Rev Respir Dis* 144:848-854 (1991).
- Adamson IYR, Bowden DH. Role of polymorphonuclear leukocytes in silica-induced pulmonary fibrosis. *Am J Pathol* 117:37-43 (1984).
- Morimoto Y, Kido M, Tanaka I, Fujino A, Higashi T, Yokosaki Y. Synergistic effects of mineral fibres and cigarette smoke on the production of tumour necrosis factor by alveolar macrophages of rats. *Br J Ind Med* 50:955-960 (1993).
- Morimoto Y, Tanaka I, Higashi T, Hori H, Yamato H, Kido M, Yokosaki Y. Chronic exposure to cigarette smoke inhibits the stimulating effects of mineral fibers on the production of tumor necrosis factor by alveolar macrophages of rats. *Int J Occup Environ Health* 2:95-98 (1996).
- Remick DG, DeForge LE. Cytokines and pulmonary fibrosis. In: *Pulmonary Fibrosis* (Phan SH, Thrall RS, eds). New York:Marcel Dekker, Inc., 1995:599-626.
- Morimoto Y, Yokosaki Y, Kido M, Tanaka I, Fujino A, Higashi T. Mineral fibers stimulate alveolar macrophages to produce tumor necrosis factor: influence of type and size. *Jpn J Traumatol Occup Med* 41:386-393 (1993).
- Forget G, Lacroix MJ, Brown RC, Evans PH. Response of perfused alveolar macrophages to glass fibers: effect of exposure duration and fiber length. *Environ Res* 39:124-35 (1986).
- Tilkes F, Beck EG. Influence of well-defined mineral fibers on proliferating cells. *Environ Health Perspect* 51:275-279 (1983).
- Zhang Y, Lee TC, Guillemin B, Yu M-C, Rom WN. Enhanced IL-1 $\beta$  and tumor necrosis factor- $\alpha$  release and messenger RNA expression in macrophages from idiopathic pulmonary fibrosis or after asbestos exposure. *J Immunol* 150:4188-4196 (1993).
- Ljungman AG, Lindahl M, Tagesson C. Asbestos fibres and man made mineral fibres: induction and release of tumour necrosis factor- $\alpha$  from rat alveolar macrophages. *Occup Environ Med* 51:777-783 (1994).
- Blackford JA, Antonini JM, Castranova V, Dey RD. Intratracheal instillation of silica up-regulates inducible nitric oxide synthase gene expression and increases nitric oxide production in alveolar macrophages and neutrophils. *Am J Respir Cell Mol Biol* 11:426-431 (1994).
- Quinlan TR, Marsh JP, Janssen YMW, Leslie KO, Hemenway D, Vacek P, Mossman BT. Dose-responsive increases in pulmonary fibrosis after inhalation of asbestos. *Am J Respir Crit Care Med* 150:200-206 (1994).
- Morimoto Y, Tsuda T, Nakamura H, Hori H, Yamato H, Nagata N, Higashi T, Kido M, Tanaka I. Expression of matrix metalloproteinases, tissue inhibitors of metalloproteinases, and extracellular matrix mRNA following exposure to mineral fibers and cigarette smoke *in vivo*. *Environ Health Perspect* 105(suppl 5):1247-1251 (1997).
- Lyons JG, Birkedal-Hansen B, Pierson MC, Whitelock JM, Birkedal-Hansen H. Interleukin-1 beta and transforming growth factor-alpha/epidermal growth factor induce expression of M(r) 95,000 type IV collagenase/gelatinase and interstitial fibroblast-type collagenase by rat mucosal keratinocytes. *J Biol Chem* 268(25):19143-19151 (1993).
- Jackson JH, Schraufstatter IU, Hyslop PA, Vosbeck K, Sauerheiser R, Weitzman SA, Cochrane CG. Role of oxidants of DNA damage; Hydroxyl radical mediates the synergistic DNA damaging effects of asbestos and cigarette smoke. *J Clin Invest* 80:1090-1095 (1987).
- Kane AB, MacDonald JL. Mechanisms of mesothelial cell injury, proliferation, and neoplasia induced by asbestos fibers. In: *Fiber Toxicology* (Warheit DB, ed). San Diego, CA:Academic Press, 1993:323-347.